



REFERENCE CHARTS OF EVIDENCE-BASED PRACTICES

ADJUSTMENT DISORDER

What Works	
There are currently no evidence-based practices.	
What Seems to Work	
Interpersonal psychotherapy (IPT)	IPT helps children and adolescents address problems to relieve depressive symptoms.
Cognitive behavioral therapy (CBT)	CBT is used to improve age-appropriate problem-solving skills, communication skills, and stress management skills. It also helps the child's emotional state and support systems to enhance adaptation and coping.
Stress management	Stress management is particularly beneficial in cases of high stress and helps the youth learn how to manage stress in a healthy way.
Group therapy	Group therapy among like-minded/afflicted individuals can help group members cope with various features of adjustment disorders.
Family therapy	Family therapy is helpful for identifying needed changes within the family system. These changes may include improving communication skills and family interactions and increasing support among family members.
What Does Not Work	
Medication alone	Medication is seldom used as a singular treatment because it does not aid the child in learning how to cope with the stressor. Targeted symptomatic treatment of the anxiety, depression, and insomnia may effectively augment therapy.

ANOREXIA NERVOSA

What Works	
Family-based psychotherapy	Family members are included in the process to assist in reduction of symptoms and modify maladaptive interpersonal patterns.
In-patient behavioral programs	Individuals are rewarded for engaging in healthy eating and weight-related behaviors.

Nutritional rehabilitation	Entails developing meal plans and monitoring intake of adequate nutrition to promote healthy weight gain.
What Seems to Work	
Cognitive behavioral therapy (CBT)	Needs further study to be well established; it is used to change underlying eating disorder cognitions and behaviors.
Medication	Used primarily after weight restoration to minimize symptoms associated with psychiatric comorbidities.
Not Adequately Tested	
Individual psychotherapy	Controlled trials have not supported this treatment; however, it may be beneficial during the renourishing process and to minimize comorbid symptoms.
Imaginal Exposure therapy	Involves imagining fear-inducing situations and then working through resulting emotions and behaviors. Future randomized control trials still needed.
What Does Not Work	
Group psychotherapy	May stimulate the transmission of unhealthy techniques among group members, particularly during acute phase of disorder.
12-step programs	Not yet tested for their efficacy; discouraged as a sole treatment.
Tricyclic antidepressants	Tricyclic antidepressants are contraindicated and should be avoided in underweight individuals and in individuals who are at risk for suicide.
Somatic treatments	To date, treatments such as vitamin and hormone treatments and electroconvulsive therapy show no therapeutic value.

ANXIETY DISORDERS

What Works	
Behavioral (e.g., exposure) & cognitive behavioral (CBT) therapies	Treatment that involves exposing youth to the (non-dangerous) feared stimuli and challenging the cognitions associated with the feared stimuli, with the goal of the youth's learning that anxiety decreases over time.
Selective serotonin reuptake inhibitors (SSRIs)	Treatment with certain SSRIs have been proven to help with anxiety; however, SSRIs may increase suicidal ideation in some youth.
What Seems to Work	
Educational support	Psychoeducational information on anxiety provided to parents, usually in a group setting.
Benzodiazepines	While proven effective, not a first choice treatment because of an increase in the risk of behavioral disinhibition.
Computer-based behavioral & cognitive behavioral therapy (CBT)	CBT administered electronically to eliminate long waiting periods or lack of clinical experts in a given area.

Not Adequately Tested	
Play therapy	Therapy using self-guided play to encourage expression of feelings and healing.
Antihistamines or herbs	No controlled studies on efficacy.
Psychodynamic therapy	Therapy designed to uncover unconscious psychological processes to alleviate the tension thought to cause distress.
Neurofeedback	A type of non-invasive brain training that enables an individual to learn how to change mental and/or physiological activity.
Antipsychotics/ neuroleptics	High level of risk of impaired cognitive functioning and tardive dyskinesia with long-term use; contraindicated in youth who do not also have Tourette's syndrome or psychosis.

ATTENTION-DEFICIT/HYPERACTIVITY DISORDER (ADHD)

What Works	
Behavioral classroom management (BCM)	BCM uses contingency management strategies, including teacher-implemented reward programs, token systems, time-out procedures, and daily report cards (DRCs). Clinicians or parents may work in consultation with teachers to develop a classroom treatment plan.
Behavioral parent training (BPT)	BPT teaches the parent to implement contingency management strategies similar to BCM techniques at home.
Intensive behavioral peer intervention (BPI)	Intensive BPI is conducted in recreational settings, such as summer treatment programs (STPs). STPs have demonstrated effectiveness and are considered well-established. However, STPs are less feasible to implement than other evidence-based practices.
Stimulant: d-Amphetamine	Short-acting: Adderall, Dexedrine, DextroStat Long-acting: Dexedrine Spansule, Adderall XR, Lisdexamfetamine
Stimulant: Methylphenidate	Short-acting: Focalin, Methylin, Ritalin; Intermediate-acting: Metadate ER, Methylin ER, Ritalin SR, Metadate CD, Ritalin LA; Long-acting: Concerta, Daytrana patch, Focalin XR
Serotonin and norepinephrine reuptake inhibitor (SNRI): atomoxetine	SNRIs do not carry the same risk for addiction as some other medications.
Non-stimulant: Alpha-agonists: guanfacine and clonidine	Can be an alternative for children who do not tolerate stimulants well; both are available in short and long-acting forms.

What Seems to Work	
Non-stimulant: Serotonin and norepinephrine reuptake inhibitor (SNRI)- viloxazine	FDA-approved in April 2021 for ADHD in youth ages 6 to 17.
Not Adequately Tested	
Dietary interventions	Interventions include elimination of food additives, elimination of allergens/sensitivities, and use of nutritional supplements. Best viewed as a potential complementary intervention.
Interactive metronome training	Involves synchronizing of hand and foot exercises to audible tones.
Neurofeedback	Involves monitoring brain waves and rewarding focused attention through computerized games and exercises.
Antidepressants	These include bupropion (Wellbutrin), imipramine (Tofranil), nortriptyline (Pamelor, Aventil).
What Does Not Work	
Cognitive, psychodynamic, and client-centered therapies	Some talk therapies and some forms of play therapy have been demonstrated to have little to no effect on ADHD symptoms. ADHD is best treated with intensive behavioral interventions in the youth’s environment.
Office-based social skills training	Once-weekly, office-based training, either one-on-one or in a group setting, has not led to significant improvement in social skills. However, intensive group social skills training that uses behavioral interventions is considered well-established.

AUTISM SPECTRUM DISORDER (ASD)

What Works	
Applied behavior analysis (ABA)	Uses principles of learning theory to bring about meaningful and positive change in behavior, build a variety of skills (e.g., communication, social skills, self-control, and self-monitoring), and help generalize these skills to other situations; also known as early intensive behavioral intervention and comprehensive behavioral treatment for young children (CBTYC).
Discrete trial teaching or training (DTT)	A behavioral intervention that uses operant learning techniques to change behavior. Also known as the ABC model (action request, behavior, consequence).
Pivotal response training (PRT)	Involves targeting pivotal behaviors related to motivation to engage in social communication, self-initiation, self-management, and responsiveness to multiple cues.

Reference Charts of Evidence-Based Practices

Positive behavioral interventions	Behavioral interventions analyze the cause of a negative behavior and how it is being reinforced, and then offer techniques targeted to promoting positive behaviors.
Modeling	Involves demonstrating a target behavior to encourage imitation.
Story-based intervention	Uses stories to increase perspective-taking skills.
Self-management	Strategies that involve teaching youth to track performance while completing an activity.
Cognitive behavioral intervention package	CBT modified for ASD youth.
Language training	Targets the ability to communicate verbally.
Scripting	Provides scripted language to be used as a model in specific situations.
Naturalistic teaching strategies (NTS)	Child-directed strategies that use naturally occurring activities to increase adaptive skills.
Peer training package	Involves training peers on how to behave during social interactions with a youth with ASD.
Schedules	Used to increase independence in youth with ASD.
Learning experience: An alternative program (LEAP)	A type of peer training program for peers, teachers, parents, and others.
Social skills package	Aims to provide youth with the skills (such as making eye contact appropriately) necessary to participate in social environments.
Parent training package	Involves training parents to act as therapists.
What Seems to Work	
Augmentative and alternative communication devices	Communication systems designed to complement speech (pictures, symbols, communication boards, or other assistive technology, like tablets, text-to-speech programs, etc.).
Developmental relationship-based treatment	Programs that emphasize the importance of building social relationships by using the principals of developmental theory.
Exercise	Uses physical exertion to regulate behavior and help with social, communication, and motor skills. The value of exercise is not that stereotypic behaviors stop, but that stress, dysregulation, or feelings of being overwhelmed decrease as a result of physical activity.
Exposure package	Involves gradually exposing youth to the non-dangerous situations that they fear, with a focus on having them learn that their anxiety will decrease over time; at the same time, the use of maladaptive strategies used in the past is prevented.

Reference Charts of Evidence-Based Practices

Functional communication training (FCT)	Behavioral method that replaces disruptive or inappropriate behavior with more appropriate and effective communication.
Imitation-based intervention	Relies on adults imitating the actions of a child; promotes a “back-and forth” dynamic and the use of spontaneous language/vocalizations.
Initiation training	Involves directly teaching individuals with ASD to initiate interactions with their peers.
Language training (production and understanding)	Aims to increase both speech production and understanding of communicative acts.
Massage therapy	Uses deep tissue stimulation to induce physiological reactions in the body that have preventive, therapeutic, and relaxing effects.
Medication	Risperidone and aripiprazole are FDA-approved to target behavioral symptoms of ASD like aggression, self-injury, and irritability.
Multi-component package	Involves a combination of multiple treatment procedures that are derived from different fields of interest or different theoretical orientations.
Music therapy	Uses music to practice communication and social skills or behavioral goals.
Picture exchange communication system	Involves an alternative communication system designed to teach functional communication to youth with limited skills.
Reductive package	Relies on strategies designed to reduce problem behaviors without increasing alternative appropriate behaviors.
Sign language instruction	Teaches sign language as a means of communicating.
Social communication intervention	Targets some combination of social communication impairments.
Structured teaching	Relies heavily on the visual organization, predictable schedules, and individualized use of teaching methods; can be used in educational, community, and home settings.
Technology-based intervention	Presents instructional materials using the medium of computers or related technologies.
Theory of mind training	Aims to teach youth to recognize and identify the mental states of others.
Not Adequately Tested	
<ul style="list-style-type: none"> • Animal-assisted therapy (e.g., hippotherapy: the use of horseback riding as a therapeutic or rehabilitative treatment) • Auditory integration training • Concept mapping • DIR/Floortime • Gluten-free and/or casein-free diet 	<ul style="list-style-type: none"> • Movement-based intervention • SENSE theatre intervention • Sensory intervention package • Social-behavioral learning strategy • Social cognition intervention • Social thinking intervention

What Does Not Work

Facilitated communication	A person with a disability is assisted by a facilitator in typing letters, words, phrases, or sentences on a keyboard.
---------------------------	--

BINGE EATING DISORDER**What Works**

There are no evidence-based practices at this time.

What Seems to Work

Cognitive behavioral therapy (CBT)	The most effective independent treatment option. It is used to change underlying eating disorder cognitions and behaviors.
Interpersonal psychotherapy (IPT)	Attempts to reduce the use of binge eating as a coping mechanism by supporting the development of healthy interpersonal skills.
Medication	Antidepressants, namely SSRIs, have effectively reduced binge/purging behaviors, as well as comorbid psychiatric symptoms.

Not Adequately Tested

Dialectical behavior therapy (DBT)	Combines parts of CBT with principles of mindfulness.
Mindfulness and yoga-based interventions	Use of yoga as a therapeutic model to incorporate concentration, meditation, and physical and emotional awareness practices.

What Does Not Work

Nutritional rehabilitation and counseling	Although initial weight loss is associated with these treatments, weight is commonly regained.
12-step programs	Discouraged as a sole treatment because they do not address nutritional or behavioral concerns.

BIPOLAR AND RELATED DISORDERS**What Works**

There are no evidence-based practices at this time.

What Seems to Work

Medication	Mood stabilizers (lithium)/Anticonvulsants, Second-generation antipsychotics
Family-focused psychoeducational therapy (FFT)	Helps youth make sense of their illness and accept it and also to better understand use of medication; also helps to manage stress, reduce negative life events, and promote a positive family environment.

Child- and family-focused cognitive behavioral therapy (CFF-CBT)	Emphasizes individual psychotherapy with youth and parents, parent training and support, and family therapy.
Multifamily psychoeducation groups (MFPG)	Youth and parent group therapy have been shown to increase parental knowledge, promote greater access to services, and increase parental social support for youth.
Not Adequately Tested	
Interpersonal social rhythm therapy (IPSRT)	Works to minimize the effects of life stressors by helping youth establish regular patterns of sleep, exercise, and social interactions.
Omega-3 fatty acids	Unclear if supplementation helps with depressive symptoms when used in conjunction with other treatments.
Topiramate Oxcarbazepine	Anticonvulsants, not proven to be effective in youth or adults.
Dialectical behavior therapy (DBT)	Family skills training and individual therapy, not proven to help with mania or interpersonal functioning.

BODY DYSMORPHIC DISORDER

What Works

There are no evidence-based practices at this time.

Not Adequately Tested

Selective serotonin reuptake inhibitors (SSRIs)	Possibly effective because of effectiveness with similar disorders.
Cognitive behavioral therapy (CBT)	Shows promise because of its effectiveness with similar disorders.

BULIMIA NERVOSA

What Works

Cognitive behavioral therapy (CBT)	The most effective independent treatment option; it is used to change underlying eating disorder cognitions and behaviors.
Combined treatments	A combination of CBT and medication seems to maximize outcomes.

What Seems to Work

Medication	Antidepressants, namely SSRIs, have effectively reduced binge/purging behaviors, as well as comorbid psychiatric symptoms.
------------	--

Not Adequately Tested	
Individual psychotherapy	Compared to CBT, few individual therapeutic approaches have been effective in reducing symptoms.
Family therapy	May be more beneficial than individual psychotherapy, but outcomes should be considered preliminary at this time.
What Does Not Work	
Bupropion	Bupropion has been associated with seizures in purging individuals with BN and is contraindicated.
Monoamine oxidase inhibitors (MAOIs)	MAOIs are potentially dangerous in individuals with chaotic bingeing and purging and their use is contraindicated.
12-step programs	Discouraged as a sole treatment because they do not address nutritional or behavioral concerns.

DEPRESSIVE DISORDERS – CHILDREN

What Works	
Stark’s cognitive behavioral therapy (CBT)	Stark’s CBT (child only or child plus parent) includes mood monitoring, mood education, increasing positive activities and positive self-statements, and problem solving. CBT is based on the premise that patients with depression have thinking that is characterized by dysfunctional negative views of oneself, experiences, and the future. CBT helps individuals identify and modify maladaptive thinking and behavior patterns.
Fluoxetine in combination with CBT	Fluoxetine, a selective serotonin reuptake inhibitor (SSRI), is the only antidepressant approved by the FDA for use in children (eight years old or older) for depression. For moderate to severe depression, fluoxetine in combination with psychosocial therapy may be warranted. However, because SSRIs can increase suicidal behavior in youth, children taking fluoxetine must be closely monitored by a mental health professional.
What Seems to Work	
Penn prevention program (PPP)	PPP is a CBT-based program that targets pre-adolescents and early adolescents who are at risk for depression.
Self-control therapy	Self-control therapy is a school-based CBT that focuses on self-monitoring, self-evaluating, and causal attributions.
Behavioral therapy	Behavioral therapy includes pleasant activity monitoring, social skills training, and relaxation.

DEPRESSIVE DISORDERS – ADOLESCENTS

What Works	
Cognitive behavioral therapy (CBT) provided in a group setting	CBT for depression focuses on identifying thought and behavior patterns that lead to or maintain the problematic symptoms.
Interpersonal therapy (IPT)	In IPT, the therapist and patient address the patient’s interpersonal communication skills, interpersonal conflicts, and family relationship problems.
Fluoxetine in combination with CBT	Fluoxetine, a selective serotonin reuptake inhibitor (SSRI), is the only antidepressant approved by the FDA for use in children (eight years old or older) for depression. For moderate to severe depression, fluoxetine in combination with psychosocial therapy may be warranted. However, because SSRIs can increase suicidal behavior in youth, children taking fluoxetine must be closely monitored by a mental health professional.
What Seems to Work	
CBT in a group or individual setting with a parent/family component	CBT for depression focuses on identifying thought and behavior patterns that lead to or maintain the problematic symptoms.
Adolescent coping with depression (CWD-A)	CWD-A includes practicing relaxation and addressing maladaptive patterns in thinking, as well as scheduling pleasant activities and learning communication and conflict resolution skills.
Interpersonal psychotherapy for depressed adolescents (IPT-A)	IPT-A addresses the adolescent’s specific interpersonal relationships and conflicts, and helps the adolescent be more effective in their relationships with others.
Physical exercise	Physical exercise has shown promise in improving symptoms of depression in adolescents. Group-based and supervised light- or moderate-intensity exercise activities three times a week for a period of between 6 to 11 or 12 weeks may bring about an improvement in depression.
Not Adequately Tested	
Dietary supplements	Supplements, such as St. John’s Wort, SAM-e, and Omega-3 have not been adequately tested and may have harmful side effects or interact with other medications. Parents should discuss supplement use with a mental health care professional.
What Does Not Work	
Tricyclic antidepressants	These antidepressants can have problematic side effects and are not recommended for children or adolescents with depression.

DEVELOPMENTAL COORDINATION DISORDER

What Works	
There are no evidence-based practices at this time.	
What Seems to Work	
Cognitive motor intervention	Therapists design a set of exercises into steps for children to practice at home. Emotional, motivational, and cognitive aspects are emphasized, as children are taught how to plan a movement, how to execute it, and how to evaluate their success. Building self-confidence through positive reinforcement is a critical goal, as success depends upon the patient’s motivation to practice outside of therapy.
Physical and occupational therapy	Tailored to a child’s specific needs.

DISRUPTIVE, IMPULSE-CONTROL, AND CONDUCT DISORDERS

What Works	
Parent management training (PMT)	<p>PMT programs focus on teaching and practicing parenting skills with parents or caregivers. Program models include:</p> <ul style="list-style-type: none"> • Helping the Noncompliant Child • Incredible Years • Parent-child interaction therapy • Parent MT to Oregon model
Multisystemic therapy (MST)	MST is an intensive family- and community- based treatment that addresses the multiple determinants of serious antisocial behavior. MST clinicians use empirically validated approaches, such as cognitive-behavioral therapy and pragmatic family therapies, and typically provide individual and family counseling and 24-hour crisis management.
Cognitive behavioral therapy (CBT)	CBT emphasizes problem-solving skills and anger control/coping strategies.
CBT & parent management training	Combines CBT and PMT.
What Seems to Work	
Multidimensional treatment foster care	Community-based program alternative to institutional, residential, and group care placements for use with severe chronic delinquent behavior; foster parents receive training and provide intensive supported treatment within the home.
Not Adequately Tested	
Brief Strategic Family Therapy	Counselors establish relationships with the family, then coach to reshape patterns of interpersonal interactions.

Atypical antipsychotics medications	Risperidone (Risperdal), quetiapine (Seroquel), olanzapine (Zyprexa), and aripiprazole (Abilify); limited evidence for effectiveness in youth with ID or ASD; primarily short-term use recommended.
Stimulant or atomoxetine	Methylphenidate, d-Amphetamine, atomoxetine; limited evidence when comorbid with primary diagnosis of ADHD.
Mood stabilizers	Divalproex sodium, lithium carbonate; limited evidence when comorbid with primary diagnosis of bipolar disorder.
Selective serotonin reuptake inhibitors (SSRIs)	Limited evidence when comorbid with primary diagnosis of depressive disorder; may also address aggression.
What Does Not Work	
Boot camps, shock, incarcerations	Ineffective at best; can lead to worsening of symptoms.
Dramatic, short-term, or talk therapy	Little to no effect as currently studied.

FIRESETTING, JUVENILE

What Works	
There are no evidence-based practices at this time.	
What Seems to Work	
Cognitive behavioral therapy (CBT)	Structured treatments designed to intervene with children who set fires. Because firesetting is a maladaptive behavior, CBT is a reasonable intervention to consider for behavior modification.
Fire safety education	Education includes information about the nature of fire, how rapidly it spreads, and its potential for destructiveness, as well as information about how to maintain a fire-safe environment, utilizing escape plans and practice, and the appropriate use of fire.
Firefighter home visit	Firefighters visit homes and explain the dangers of playing with fire.
What Does Not Work	
Ignoring the problem	Leaving youth untreated is not beneficial because they typically do not outgrow this behavior. Behavior may increase.
Satiation	Satiation, repetitively lighting and extinguishing fires, may cause the youth to feel more competent around fire and may actually increase the behavior.
Burning the juvenile	Burning a juvenile to show the destructive force of fire is illegal and abusive. It will not decrease the likelihood of the juvenile setting fires or treat the problem.

Scaring the juvenile	Scare tactics may produce the emotions or stimulate the actions the clinician is trying to prevent, particularly when family or social issues may trigger firesetting. Scare tactics may also trigger defiance, avoidance, or may even increase the likelihood that firesetting traits continue.
----------------------	--

HOARDING DISORDER

What Works	
There are no evidence-based practices at this time.	
What Seems to Work	
Cognitive behavioral therapy (CBT) for hoarding	A multi-component, cognitive behavioral treatment designed specifically for hoarding has shown promising results in adults.
Not Adequately Tested	
Selective serotonin reuptake inhibitors (SSRIs)	Possibly effective because of their effectiveness with similar disorders.

INTELLECTUAL DISABILITY (ID)

What Works	
Behavioral interventions	Behavioral interventions analyze the cause of a negative behavior and how it is being reinforced and then offer techniques targeted to promoting positive behaviors. They may also involve modified teaching approaches to assist in the development of social skills, emotion regulation, and functional skills.
Applied behavioral analysis (ABA)	A type of behavioral intervention that uses principles of learning theory to bring about meaningful and positive change in behavior. ABA techniques have been used to help build a variety of skills (e.g., communication, social skills, self-control, and self-monitoring) and help generalize these skills to other situations.
Functional communication training (FCT)	An example of a behavioral intervention program that combines the assessment of the communicative functions of problem behavior with ABA procedures to teach alternative responses. Problem behaviors can be eliminated through extinction and replaced with alternate, more appropriate forms of communicating needs or wants.
What Seems to Work	
Psychotropic medications for co-occurring mental health disorders	Prescribed to treat co-occurring disorders such as anxiety disorders and ADHD. Because these medications have not been studied in ID populations, they should only be used when therapeutic and social measures do not properly address symptoms and in conjunction with appropriate behavioral interventions.

Not Adequately Tested

Psychotropic medications to treat challenging behaviors	Psychotropic medications are sometimes used “off label” to treat challenging behaviors such as aggression. These medications should be used with caution and only when necessary. They should never be used for the convenience of caregivers.
---	--

JUVENILE OFFENDING

What Works

Multisystemic therapy (MST)	An integrative, family-based treatment with a focus on improving psychosocial functioning for youth and families.
Functional family therapy (FFT)	A family-based program that focuses on delinquency, treating maladaptive and “acting out” behaviors, and identifying obtainable changes.
Treatment Foster Care Oregon (TFCO)	As an alternative to corrections or residential treatment, TFCO places juvenile offenders with carefully trained foster families who provide youth with close supervision, fair and consistent limits, consequences, and a supportive relationship with an adult. The program includes family therapy for biological parents, skills training and supportive therapy for youth, and school-based behavioral interventions and academic support.

What Seems to Work

Family centered treatment (FCT)	FCT seeks to address the causes of parental system breakdown while integrating behavioral change. FCT provides intensive in-home services and is structured into four phases: joining and assessment, restructuring, value change, and generalization.
Brief strategic family therapy	A short-term, family-focused therapy that focuses on changing family interactions and contextual factors that lead to behavior problems.
Aggression replacement therapy (ART)	A short-term, educational program that focuses on anger management and provides youth with the skills to demonstrate non-aggressive behaviors, decrease antisocial behaviors, and utilize prosocial behaviors.
Cognitive behavioral therapy (CBT)	A structured, therapeutic approach that involves teaching youth about the thought-behavior link and working with them to modify their thinking patterns in a way that will lead to more adaptive behavior in challenging situations.
Dialectical behavior therapy (DBT)	A therapeutic approach that includes individual and group therapy components and specifically aims to increase self-esteem and decrease self-injurious behaviors and behaviors that interfere with therapy.

NONSUICIDAL SELF-INJURY

What Works	
There are no evidence-based practices at this time.	
What Seems to Work	
Cognitive behavioral therapy (CBT)	CBT involves providing skills designed to assist youth with affect regulation and problem solving.
Dialectical behavior therapy (DBT)	DBT emphasizes acceptance strategies and the development of coping skills.
Not Adequately Tested	
Problem solving therapy	Designed to improve an individual’s ability to cope with stressful life experiences.
Medication	Evidence of the effectiveness of the use of medications, such as high-dose SSRIs, atypical neuroleptics, and opiate antagonists, is limited. In addition, some medications have been shown to increase suicidal ideation in children and adolescents.
Hospitalization	Because effectiveness is not consistently demonstrated, should be reserved for youth who express intent to die.
What Does Not Work	
Benzodiazepines or trazodone	May potentially increase incidence of NSSIs in depressed or anxious individuals.

OBSESSIVE-COMPULSIVE DISORDER (OCD)

What Works	
Cognitive behavioral therapy (CBT) with exposure and response prevention (ERP)	Treatment path with a consistent and compelling relationship between the disorder, the treatment, and the specified outcome; combines training with exposure and preventing the accompanying response.
Family-focused individual CBT	Individual CBT that includes a focus on family involvement. It should be noted that the distinction of family focused here is meant to imply a format for treatment delivery.
Serotonin reuptake inhibitors (SRIs)	Clomipramine: Approved for children aged ten and older; recommend periodic electrocardiographic (ECG) monitoring.
Selective serotonin reuptake inhibitors (SSRIs)	Fluoxetine (Prozac): Approved for children aged eight and older. Sertraline (Zoloft): Approved for children aged six and older. Fluvoxamine (Luvox): Approved for children aged eight and older.

What Seems to Work	
Family-focused group CBT	Studies show promising results, but there have only been a small number of studies. However, each study addresses complex comorbidity and issues impacting community-based treatment.
Not Adequately Tested	
CBT without ERP Psychodynamic therapy Client-centered therapy	Systematic controlled studies have not been conducted using these approaches.
Technology-based CBT	Results show preliminary support for telephone CBT and web-camera CBT. Although these results are encouraging, caution must be taken due to the small sample sizes and lack of active control groups.
Deep Brain Stimulation (DBS)	Involves implanting electrodes in the brain to monitor abnormal impulses. There is strong evidence for success in young adults, but not enough adolescent studies.
What Does Not Work	
Antibiotic treatments	Antibiotic treatments are only indicated when the presence of an autoimmune or strep-infection has been confirmed and coincided with onset or increased severity of obsessive-compulsive disorder symptoms (PANDAS).
Herbal therapies	Herbs, such as St. John’s Wort, have not been rigorously tested and are not FDA approved. In some instances, herbal remedies may make symptoms worse or interfere with medications.

POST-TRAUMATIC STRESS DISORDER (PTSD)

What Works	
Trauma-focused cognitive behavioral therapy (TF-CBT)	Treatment that involves reducing negative emotional and behavioral responses related to trauma by providing psychoeducation on trauma, addressing distorted beliefs and attributes related to trauma, introducing relaxation and stress management techniques, and developing a trauma narrative in a supportive environment.
What Seems to Work	
Family centered treatment (FCT) trauma treatment	FCT trauma treatment provides intensive in-home services and seeks to address the causes of trauma, including parental system breakdown, while integrating behavioral change.
School-based group CBT	Similar components to TF-CBT, but in a group, school-based format.
Not Adequately Tested	
Child-centered play therapy	Therapy that utilizes child-centered play to encourage expression of feelings and healing.
Medication	Includes treatment with selective serotonin reuptake inhibitors (SSRIs).
Resilient peer treatment	Classroom treatment that pairs withdrawn children with resilient peers with a parent present for assistance.

Eye movement desensitization and reprocessing therapy (EMDR)	Therapy that utilizes visual and physical memory imagery while the clinician creates visual or auditory stimulus to reduce negative memory and increase positive memory.
What Does Not Work	
Restrictive rebirthing or holding techniques	Restrictive rebirthing or holding techniques that may forcibly bind or restrict, coerce, or withhold food/water from children and have resulted, in some cases, in death; not recommended.
Psychological debriefing	An approach in which youth talk about the facts of the trauma (and associated thoughts and feelings) and then are encouraged to re-enter into the present. Recent research suggests this approach is ineffective and potentially harmful.

SCHIZOPHRENIA

What Works	
There are no evidence-based practices at this time.	
What Seems to Work	
Medication treatment with second-generation (atypical) antipsychotics	Risperidone Aripiprazole Quetiapine Paliperidone Olanzapine
Medication treatment with traditional neuroleptics/first generation antipsychotics	Molindone Haloperidol
Family psychoeducation and support	Helps to improve family functioning, problem solving, and communication skills, and decreases relapse rates.
Cognitive behavioral therapy (CBT)	Includes social skills training, problem-solving strategies, and self-help skills.
Cognitive remediation	Pointed tasks to help improve specific deficiencies in cognitive, emotional, or social aspects of a patient’s life.
Not Adequately Tested	
Electroconvulsive therapy (ECT)	Small electric currents are passed through the brain, intentionally triggering a brief seizure to reverse symptoms of certain mental illnesses. Unproven as effective in youth. Should only be used as a last effort after all risks are weighted against possible benefits.
What Does Not Work	

Psychodynamic therapies	Talk therapies that focus on a client's self-awareness and understanding of the influence of the past on present behavior. These therapies are considered to be potentially harmful for youth with schizophrenia.
-------------------------	---

SEXUAL OFFENDING

What Works	
There are no evidence-based practices at this time.	
What Seems to Work	
Multisystemic therapy for problem sexual behaviors (MST-PSB)	An intensive family and community based treatment that addresses the multiple factors of serious antisocial behavior in juvenile sexual abusers.
Cognitive behavioral therapy (CBT); Children with problematic sexual behavior CBT (PBS-CBT)	Treatment modalities that provide cognitive-behavioral, psychoeducational, and supportive services.
Not Adequately Tested	
Medication	There is no research validation for the use of medication targeting sexually deviant behavior in youth and only limited methodologically sound research to guide in the treatment of adults.

STEREOTYPIC MOVEMENT DISORDER

What Works	
Habit reversal therapy (HRT)	Increases awareness to the feelings and context associated with the stereotypes and implements competing and inconspicuous habits in their place. HRT can be modified to include rewards, relaxation, education, self-awareness, and situational changes. It is sometimes combined with other therapies.
What Seems to Work	
Medication	Medications may be considered for moderate to severe stereotypes causing severe impairment in quality of life or when co-occurring conditions that would also benefit from the medication are present.

SUBSTANCE USE DISORDERS

What Works	
Cognitive behavioral therapy (CBT)	A structured therapeutic approach that involves teaching youth about the thought-behavior link and working with them to modify their thinking patterns in a way that will lead to more adaptive behavior in challenging situations.

Family therapy Multidimensional family therapy (MDFT) Functional family therapy (FFT)	Family-based therapy is aimed at providing education, improving communication and functioning among family members, and reestablishing parental influence through parent management training. MDFT views drug use in terms of networks of influences (individual, family, peer, community) and encourages treatment across settings in multiple ways. FFT is best used in youth with co-occurring conduct and delinquent behaviors. It combines a focus on family relationships with CBT interventions to change patterns and improve the family's functioning.
Multisystemic therapy (MST)	An integrative, family-based treatment with a focus on improving psychosocial functioning for youth and families.
What Seems to Work	
Behavioral therapies	Behavioral therapies focus on identifying specific problems and areas of deficit and working on improving these behaviors.
Motivational interviewing (MI) Motivational enhancement therapy (MET)	MI is a brief treatment approach aimed at increasing motivation for behavior change. It is focused on expressing empathy, avoiding argumentation, rolling with resistance, and supporting self-efficacy. MET is an adaptation of MI that includes one or more client feedback sessions in which normative feedback is presented and discussed.
Medication	Some medication can be used for detoxification purposes, as directed by a doctor. Medication may also be used to treat co-existing mental health disorders.
Not Adequately Tested	
Multifamily educational intervention (MEI)	MEI combines psycho-educational and family interventions for troubled adolescents and their families.
Adolescent group therapy (AGT)	The AGT intervention incorporates adolescent therapy groups on stress management, developing social skills, and building group social support.
Interpersonal and psychodynamic therapies	Interpersonal and psychodynamic therapies are methods of individual counseling that are often incorporated into the treatment plan and focus on unconscious psychological conflicts, distortions, and faulty learning.
Client-centered therapies	A type of therapy focused on creating a non-judgmental environment, such that the therapist provides empathy and unconditional positive regard. This facilitates change and solution making on behalf of the youth.
Psychoeducation	Programs aimed at educating youth on substance use; may cover topics like peer pressure and consequences of substance use.
Project CARE	A program aimed at raising awareness about chemical dependency among youth through education and training.
Twelve-step programs	A twelve-step program that uses the steps of Alcoholics Anonymous as principles for recovery and treating addictive behaviors.
Process groups	A type of psychotherapy that is conducted in a small group setting. Groups can be specialized for specific purposes, and therapy utilizes the group as a mechanism of change.

Neurofeedback	A type of non-invasive brain training that enables an individual to learn how to change mental and/or physiological activity.
---------------	---

SUICIDE, YOUTH

What Works	
There are no evidence-based practices at this time.	
What Seems to Work	
Cognitive behavioral therapy (CBT) Dialectical behavior therapy (DBT)	Both of these psychotherapies have shown promise in reducing suicidal ideation in some youth when paired with appropriate medication therapy. Other psychotherapies, such as interpersonal therapy for adolescents, psychodynamic therapy, and family therapy, may also be effective. Suicide focused treatment can be effective without medication therapy in some cases.
Selective serotonin reuptake inhibitors (SSRIs)	These antidepressants may help reduce suicidal ideation in teens with diagnosed depression; however, in some individuals they may cause suicidal ideation. Youth taking SSRIs must be closely monitored. Limitations: SSRIs must be taken consistently, require a therapeutic dose per individual, and can take up to 3 months to show effectiveness.
SOS (signs of suicide) prevention program	A school-based education and screening program that teaches students to recognize warning signs of depression and suicidality in themselves or their peers. Includes a screening component.
Safety Planning	Clinicians help patients identify effective coping techniques to use during crisis events (American Academy of Pediatrics).
Sources of Strength	Peer leaders model and encourage friends to name and engage trusted adults; reinforce and create an expectancy that friends ask adults for help for suicidal friends; and identify and use interpersonal and formal coping resources.
Attachment-Based Family Therapy	Most effective for youth aged 12-24; based on the idea that the quality of familial relationships has the potential to exacerbate or mitigate depression, suicide, and suicidal ideation; the only treatment adapted specifically for LGBTQ+ youth.
Not Adequately Tested	
Collaborative Assessment and Management of Suicidality (CAMS)	Centered on building a relationship between the physician and individual that allows the individual to identify and manage suicidal thoughts, feelings, and behaviors.
Gatekeeper training	Involves educating youth, parents, and caregivers about warning signs of suicide to encourage early intervention.
What Does Not Work	
Group counseling	Multiple randomized control trials fail to demonstrate efficacy.

Tricyclic antidepressants	Not recommended; effectiveness has not been demonstrated. Older tricyclic antidepressants are lethal in overdose quantities.
No-suicide contracts	Designed as an assessment tool, not a prevention tool; studies on effectiveness in reducing suicide are inconclusive and their use is discouraged, as they may be interpreted as being coercive or may encourage suicide in some individuals.

TIC DISORDERS

What Works	
Habit reversal therapy (HRT) for tic disorder	A type of cognitive behavioral therapy, HRT increases awareness to the feelings and context associated with the urge to tic, then replaces it with competing, inconspicuous habits.
Comprehensive behavioral intervention for tics (C-BIT)	Combines HRT and other approaches like education, awareness via self-monitoring, relaxation techniques, and sometimes situational changes.
What Seems to Work	
Exposure with response prevention (ERP)	Consists of repeated, prolonged exposures to stimuli that elicit discomfort and instructions to refrain from any behavior that serves to reduce discomfort.
Medication	Medications may be considered for moderate to severe tics causing severe impairment in quality of life or when co-occurring conditions that would also benefit from the medication are present.
Massed negative practice (MNP)	Treatment involves developing reactive inhibition through the child's over-rehearsal of target tic in high-risk situations.
Not Adequately Tested	
Web-based exposure with response prevention (ERP)	ERP virtual treatment is considered generally effective, but there are limited studies on how it compares with in-person treatment.
What Does Not Work	
Deep brain stimulation	A neurostimulator is surgically implanted into the brain; not proven to be effective and not recommended.
Repetitive transcranial magnetic stimulation (rTMS)	Uses magnetic fields and electrical pulses to affect neurons in the brain; safety in youth has not been established; not recommended.
Plasma exchange; Intravenous immunoglobulin (IVIG) treatment	Blood transfusions alter levels of plasma or immunoglobulin; while several of these treatments have been shown to be promising, they are not empirically supported and not recommended.

Dietary supplements (magnesium and vitamin B6); special diets	Supplements may have the potential to negatively interact with other medications; not recommended until safety in children is established.
---	--

TRICHOTILLOMANIA (HAIR PULLING) AND EXCORIATION (SKIN PICKING) DISORDER

What Works

There are no evidence-based practices at this time.

What Seems to Work

Habit reversal therapy (HRT)	Treatment increases awareness to the feelings and context associated with the urges and implements a competing and inconspicuous habit in place of the hair pulling and skin picking.
Cognitive behavioral therapy (CBT)	Treatment involves exposing children to the stimuli associated with the urge, while challenging thoughts associated with high-risk situations.

Not Adequately Tested

Selective serotonin reuptake inhibitors (SSRIs) N-acetylcysteine Naltrexone	Some demonstrated improvement on certain measures of picking behavior has been shown in some pharmacological studies of adults.
---	---

***The Collection of Evidence-based Practices for Children and Adolescents with
Mental Health Treatment Needs, 9th Edition***
Virginia Commission on Youth, 2023

The information contained herein is strictly for informational and educational purposes only and is not designed to replace the advice and counsel of a physician, mental health provider, or other medical professional. If you require such advice or counsel, you should seek the services of a licensed mental health provider, physician, or other medical professional. The Virginia Commission on Youth is not rendering professional advice and makes no representations regarding the suitability of the information contained herein for any purpose.